

Abbreviated Prescribing Information

Metoprolol, Telmisartan Tablet

METOLAR TL

METOLAR TL tablets are available as extended-release film coated tablets for oral use.

COMPOSITION

Each film coated tablet contains Metoprolol succinate equivalent to Metoprolol tartrate (as extended release) 25 mg or 50 mg, Telmisartan 40mg.

INDICATIONS AND USAGE

METOLAR TL is indicated for the treatment of essential Hypertension.

DOSAGE AND METHOD OF ADMINISTRATION

The recommended dose is 1 tablet once daily or as directed by the Physician. To be swallowed as whole with liquid and must not be chewed or crushed.

CONTRAINDICATION

This tablet is contraindicated in patients with known hypersensitivity to any of the active substance or to any of the excipients. Metoprolol is contraindicated in patients with severe hypotension, bradycardia or heart block.

SPECIAL WARNING & PRECAUTIONS FOR USE

When pregnancy is detected, discontinue Telmisartan/Metoprolol tablets as soon as possible. Patients should not interrupt or discontinue metoprolol succinate extended-release tablets without consulting the physician.

ADVERSE REACTIONS

The most common adverse reactions seen are headache, dizziness, nausea, fatigue, hypotension and bradycardia.

Abbreviated Prescribing Information

Metoprolol, Cilnidipine & Telmisartan Tablet

METOLAR TRIO

METOLAR TRIO tablets are available as extended release film coated tablet for oral use.

COMPOSITION

Each film coated tablet contains Metoprolol succinate equivalent to Metoprolol tartrate (as extended release) 25 mg or 50 mg, Telmisartan 40mg and Cilnidipine 10mg.

INDICATIONS AND USAGE

METOLAR TRIO is indicated for the treatment of patients with uncontrolled essential hypertension and stable ischemic heart disease.

DOSAGE AND METHOD OF ADMINISTRATION

The recommended dose is 1 tablet once daily or as directed by the Physician. To be swallowed as whole with liquid and must not be chewed or crushed.

CONTRAINDICATION

This tablet is contraindicated in patients with known hypersensitivity to any of the active substance or to any of the excipients. Metoprolol is contraindicated in patients with severe hypotension, bradycardia or heart block. Cilnidipine is contraindicated in unstable angina

SPECIAL WARNING & PRECAUTIONS FOR USE

When pregnancy is detected, discontinue Telmisartan/Metoprolol tablets as soon as possible. Patients should not interrupt or discontinue metoprolol succinate extended-release tablets without consulting the physician. To be used with caution in patients with hepatic dysfunction.

ADVERSE REACTIONS

The most common adverse reactions seen are headache, dizziness, nausea, fatigue. hypotension and bradycardia.

Abbreviated Prescribing Information

Metoprolol, Chlorthalidone & Telmisartan tablets

METOLAR 3D

METOLAR 3D tablets are available as extended-release film coated tablet for oral use.

COMPOSITION

Each film coated tablet contains Metoprolol succinate equivalent to Metoprolol tartrate (as extended release) 25 mg or 50 mg, Telmisartan 40mg and Chlorthalidone 12.5mg.

INDICATIONS AND USAGE

METOLAR 3D is indicated for the treatment of essential hypertension with stable CAD.

DOSAGE AND METHOD OF ADMINISTRATION

The recommended dose is 1 tablet once daily or as directed by the Physician. To be swallowed as whole with liquid and must not be chewed or crushed.

CONTRAINDICATION

This tablet is contraindicated in patients with known hypersensitivity to any of the active substance or to any of the excipients. Metoprolol is contraindicated in patients with severe hypotension, bradycardia or heart block.

SPECIAL WARNING & PRECAUTIONS FOR USE

When pregnancy is detected, discontinue Telmisartan/Metoprolol tablets as soon as possible. Hypokalemia may develop with chlorthalidone. Patients should not interrupt or discontinue metoprolol succinate extended-release tablets without consulting the physician. To be used with caution in patients with hepatic dysfunction.

ADVERSE REACTIONS

The most common adverse reactions seen are headache, dizziness, nausea, fatigue. Hypotension and bradycardia.